

Fig. 6


nsos

НеГа


257RDB


181RDB


MCF-Adr



t¥ F












Fig. 10


daunorubicin:

daunorubicin: 40 ng/ml Control: No daunorubicin 30

Fig. 11









Otsidadist otsidadist IN BONDE 10000 1000000 10000000000 100000000 Im/u19

Fig. 15









Fig. 18A

5 pfu Xvir03 20 pfu Ad312 not infected

Fig. 18B

after incubation with 5µM Irinotecan for U373 cells infected with 10 pfu dl520/cell


24 hours Fig. 19



Southern Blot Analysis

U373 cells infected with 10 pfu dl520/cell


increasing concentrations of Trichostatin after incubation for 24 hours with Fig. 20



Southern Blot Analysis


Fig. 21

after incubation with Trichostatin for 24 hours FACS analysis: CAR expression of U373 cells



Oncolytic effect of dl520 in U373 cells in combination with Irinotecan and Trichostatin

Fig. 22



BfrI – cleavage site = 3534 bp 3° UTR = 3507 - 4107 bp ORF = 2019 - 3506 bpE1B55k region: Fig. 23




Fig. 24

gtectgecegeaactetactaecttgaectaegagaeegtgtetggaaegeeg ggattgtgactgactttgctttcctgagcccgcttgcaagcagtgcagcttcccgt ccgggaacttaatgtcgtttctcagcagctgttggatctgcgccagcaggtttctg ttggagactgcagcctccgccgcttcagccgctgcagccaccgccgcg gtgggggtcttatgtattttgtatctgttttgcagcagccgccgccatgagc accaactcgtttgatggaagcattgtgagctcatatttgacaacgcgcatgcccc catgggccggggtgcgtcagaatgtgatgggctccagcattgatggtcgccc tcatccgcccgcgatgacaagttgacggctcttttggcacaattggattctttgac 5'tgaggtactgaaatgtgtgggcgtggcttaagggtgggaaagaatatatag actetgtttggatttggatcaagcaagtgtettgetgtetttatttaggggttttge